arXiv:1210.8090v1 [math.AP] 30 Oct 2012

L^p SPECTRAL MULTIPLIERS ON THE FREE GROUP $N_{3,2}$

ALESSIO MARTINI AND DETLEF MÜLLER

Abstract. Let L be the homogeneous sublaplacian on the 6-dimensional free 2-step nilpotent Lie group $N_{3,2}$ on 3 generators. We prove a theorem of Mihlin-Hörmander type for the functional calculus of L, where the order of differentiability $s > 6/2$ is required on the multiplier.

1. Introduction

The free 2-step nilpotent Lie group $N_{3,2}$ on 3 generators is the simply connected, connected nilpotent Lie group defined by the relations

$$[X_1, X_2] = Y_3, \quad [X_2, X_3] = Y_1, \quad [X_3, X_1] = Y_2,$$

where $X_1, X_2, X_3, Y_1, Y_2, Y_3$ is a basis of its Lie algebra (that is, the Lie algebra of the left-invariant vector fields on $N_{3,2}$). In exponential coordinates, $N_{3,2}$ can be identified with $\mathbb{R}^3_x \times \mathbb{R}^3_y$, where the group law is given by

$$(x, y) \cdot (x', y') = (x + x', y + y' + x \wedge x'/2)$$

and $x \wedge x'$ denotes the usual vector product of $x, x' \in \mathbb{R}^3$. The family $(\delta_t)_{t>0}$ of automorphic dilations of $N_{3,2}$, defined by

$$\delta_t(x, y) = (tx, t^2 y),$$

turns $N_{3,2}$ into a stratified group of homogeneous dimension $Q = 9$.

Let $L = -(X_1^2 + X_2^2 + X_3^2)$ be the homogeneous sublaplacian on $N_{3,2}$. L is a self-adjoint operator on $L^2(N_{3,2})$, hence a functional calculus for L is defined via spectral integration and, for all Borel functions $F : \mathbb{R} \to \mathbb{C}$, the operator $F(L)$ is bounded on $L^2(N_{3,2})$ whenever the "spectral multiplier" F is a bounded function. Here we are interested in giving a sufficient condition for the L^p-boundedness (for $p \neq 2$) of the operator $F(L)$, in terms of smoothness properties of the multiplier F.

Let $W_2^s(\mathbb{R})$ denote the L^2 Sobolev space of (fractional) order s. Then our main result reads as follows.

Theorem 1. *Suppose that a function $F : \mathbb{R} \to \mathbb{C}$ satisfies*

$$\sup_{t>0} \|\eta\, F(t\cdot)\|_{W_2^s} < \infty$$

for some $s > 6/2$ and some nonzero $\eta \in C_c^\infty(]0, \infty[)$. Then the operator $F(L)$ is of weak type $(1, 1)$ and bounded on $L^p(N_{3,2})$ for all $p \in]1, \infty[$.

Observe that the general multiplier theorem for homogeneous sublaplacians on stratified Lie groups by Christ [3] and Mauceri and Meda [16] requires the stronger regularity condition $s > Q/2 = 9/2$. To the best of our knowledge, in the case of $N_{3,2}$ none of the results and techniques known so far allowed one to go below the condition $s > Q/2$. Our result pushes the regularity assumption down to

---

2010 *Mathematics Subject Classification.* 43A85, 42B15.

Key words and phrases. nilpotent Lie groups, spectral multipliers, sublaplacians, Mihlin-Hörmander multipliers, singular integral operators.

The first-named author gratefully acknowledges the support of the Alexander von Humboldt Foundation.

$s > d/2 = 6/2$, where $d = 6$ is the topological dimension of $N_{3,2}$. We conjecture that this condition is sharp.

The problem of L^p-boundedness for spectral multipliers on nilpotent Lie groups has a long history, and the theorem by Christ and Mauceri and Meda is itself an improvement of a series of previous results (see, e.g., [4, 8, 5]). Nevertheless it is still an open question, whether the homogeneous dimension in the smoothness condition may always be replaced by the topological dimension.

It has been known for a long time [10, 17] that such an improvement of the multiplier theorem holds true in the case of the Heisenberg and related groups (more precisely, for direct products of Métivier and abelian groups; see also [11, 14]). This class of groups, however, does not include $N_{3,2}$, nor any free 2-step nilpotent group $N_{n,2}$ on n generators (see [20, §3] for a definition), except for the smallest one, $N_{2,2}$, which is the 3-dimensional Heisenberg group. The free groups $N_{n,2}$ have in a sense the maximal structural complexity among 2-step groups, since every 2-step nilpotent Lie group is a quotient of a free one. Our result should then hopefully shed some new light and contribute to the understanding of the problem for general 2-step nilpotent Lie groups.

2. Strategy of the proof

The sublaplacian L is a left-invariant operator on $N_{3,2}$, hence any operator of the form $F(L)$ is left-invariant too. Let $\mathcal{K}_{F(L)}$ then denote the convolution kernel of $F(L)$. As shown, e.g., in [14, Theorem 4.6], the previous Theorem 1 is a consequence of the following L^1-estimate.

Proposition 2. *For all $s > 6/2$, for all compact sets $K \subseteq]0,\infty[$, and for all functions $F : \mathbb{R} \to \mathbb{C}$ such that* $\operatorname{supp} F \subseteq K$,

$$\| \mathcal{K}_{F(L)} \|_1 \leq C_{K,s} \|F\|_{W_2^s}. \tag{1}$$

Let $|\cdot|_\delta$ be any δ_t-homogeneous norm on $N_{3,2}$; take, e.g., $|(x,y)|_\delta = |x| + |y|^{1/2}$. The crucial estimate in the proof of [16] of the general theorem for stratified groups, that is,

$$\|(1 + |\cdot|_\delta)^\alpha \, \mathcal{K}_{F(L)} \|_2 \leq C_{K,\alpha,\beta} \|F\|_{W_2^\beta} \tag{2}$$

for all $\alpha \geq 0$ and $\beta > \alpha$, implies (1) when $s > 9/2$, by Hölder's inequality. In order to push the condition down to $s > 6/2$, here we prove an enhanced version of (2), that is,

$$\|(1 + |\cdot|_\delta)^\alpha \, w^r \, \mathcal{K}_{F(L)} \|_2 \leq C_{K,\alpha,\beta,r} \|F\|_{W_2^\beta}, \tag{3}$$

for some "extra weight" function w on $N_{3,2}$, and suitable constraints on the exponents α, β, r.

A similar approach is adopted in the mentioned works on the Heisenberg and related groups. However, in [17] the extra weight w is the full weight $1 + |\cdot|_\delta$, while [10] employs the weight $w(x,y) = 1 + |x|$. Here instead the weight $w(x,y) = 1 + |y|$ is used, and (3) is proved under the conditions $\alpha \geq 0$, $0 \leq r < 3/2$, $\beta > \alpha + r$ (see Proposition 9 below).

The proof of (3) when $\alpha = 0$ is based on a careful analysis exploiting identities for Laguerre polynomials, somehow in the spirit of [4, 17, 19], but with additional complexity due, inter alia, to the simultanous use of generalized Laguerre polynomials of different types. The estimate for arbitrary α is then recovered by interpolation with (2). An analogous strategy is followed in [15], where identities for Hermite polynomials are used in order to prove a sharp spectral multiplier theorem for Grushin operators.

3. A JOINT FUNCTIONAL CALCULUS

It is convenient for us to embed the functional calculus for the sublaplacian L in a larger functional calculus for a system of commuting left-invariant differential operators on $N_{3,2}$. Specifically, the operators

$$L, -iY_1, -iY_2, -iY_3 \tag{4}$$

are essentially self-adjoint and commute strongly, hence they admit a joint functional calculus (see, e.g., [13]).

If $\mathbf{Y}$ denotes the "vector of operators" $(-iY_1, -iY_2, -iY_3)$, then we can express the convolution kernel $\mathcal{K}_{G(L,\mathbf{Y})}$ of the operator $G(L,\mathbf{Y})$ in terms of Laguerre functions (cf. [7]). Namely, for all $n, k \in \mathbb{N}$, let

$$L_n^{(k)}(u) = \frac{u^{-k}e^u}{n!}\left(\frac{d}{du}\right)^n (u^{k+n}e^{-u})$$

be the n-th Laguerre polynomial of type k, and define

$$\mathcal{L}_n^{(k)}(t) = 2(-1)^n e^{-t} L_n^{(k)}(2t).$$

Further, for all $\eta \in \mathbb{R}^3 \setminus \{0\}$ and $\xi \in \mathbb{R}^3$, define $\xi_\parallel^\eta$ and $\xi_\perp^\eta$ by

$$\xi_\parallel^\eta = \langle \xi, \eta/|\eta| \rangle, \qquad \xi_\perp^\eta = \xi - \xi_\parallel^\eta \eta/|\eta|.$$

Proposition 3. *Let $G : \mathbb{R}^4 \to \mathbb{C}$ be in the Schwartz class, and set*

$$m(n,\mu,\eta) = G((2n+1)|\eta| + \mu^2, \eta), \tag{5}$$

for all $n \in \mathbb{N}$, $\mu \in \mathbb{R}$, $\xi, \eta \in \mathbb{R}^3$ with $\eta \neq 0$. Then

$$\mathcal{K}_{G(L,\mathbf{Y})}(x,y) = \frac{1}{(2\pi)^6}\int_{\mathbb{R}^3}\int_{\mathbb{R}^3}\sum_{n\in\mathbb{N}} m(n,\xi_\parallel^\eta,\eta)\,\mathcal{L}_n^{(0)}(|\xi_\perp^\eta|^2/|\eta|)\,e^{i\langle \xi,x\rangle}\,e^{i\langle \eta,y\rangle}\,d\xi\,d\eta.$$

Proof. For all $\eta \in \mathbb{R}^3 \setminus \{0\}$, choose a unit vector $E_\eta \in \eta^\perp$, and set $\bar{E}_\eta = (\eta/|\eta|) \wedge E_\eta$; moreover, for all $x \in \mathbb{R}^3$, denote by x_1^η, x_2^η, $x_\parallel^\eta$ the components of x with respect to the positive orthonormal basis E^η, $\bar{E}^\eta$, $\eta/|\eta|$ of $\mathbb{R}^3$.

For all $\eta \in \mathbb{R}^3 \setminus \{0\}$ and all $\mu \in \mathbb{R}$, an irreducible unitary representation $\pi_{\eta,\mu}$ of $N_{3,2}$ on $L^2(\mathbb{R})$ is defined by

$$\pi_{\eta,\mu}(x,y)\phi(u) = e^{i\langle \eta,y\rangle} e^{i|\eta|(u+x_1^\eta/2)x_2^\eta} e^{i\mu x_\parallel^\eta}\phi(x_1^\eta + u)$$

for all $(x,y) \in N_{3,2}$, $u \in \mathbb{R}$, $\phi \in L^2(\mathbb{R})$. Following, e.g., [1, §2], one can see that these representations are sufficient to write the Plancherel formula for the group Fourier transform of $N_{3,2}$, and the corresponding Fourier inversion formula:

$$f(x,y) = (2\pi)^{-5}\int_{\mathbb{R}^3\setminus\{0\}}\int_{\mathbb{R}} \operatorname{tr}(\pi_{\eta,\mu}(x,y)\,\pi_{\eta,\mu}(f))\,|\eta|\,d\mu\,d\eta \tag{6}$$

for all $f : N_{3,2} \to \mathbb{C}$ in the Schwartz class and all $(x,y) \in N_{3,2}$, where $\pi_{\eta,\mu}(f) = \int_{N_{3,2}} f(z)\,\pi_{\eta,\mu}(z^{-1})\,dz$.

Fix $\eta \in \mathbb{R}^3 \setminus \{0\}$ and $\mu \in \mathbb{R}$. The operators (4) are represented in $\pi_{\eta,\mu}$ as

$$d\pi_{\eta,\mu}(L) = -\partial_u^2 + |\eta|^2 u^2 + \mu^2, \qquad d\pi_{\eta,\mu}(-iY_j) = \eta_j. \tag{7}$$

If h_n is the n-th Hermite function, that is,

$$h_n(t) = (-1)^n (n!\,2^n\sqrt{\pi})^{-1/2} e^{t^2/2}\left(\frac{d}{dt}\right)^n e^{-t^2},$$

and $\tilde{h}_{\eta,n}$ is defined by

$$\tilde{h}_{\eta,n}(u) = |\eta|^{1/4} h_n(|\eta|^{1/2}u),$$

then $\{\tilde{h}_{\eta,n}\}_{n\in\mathbb{N}}$ is a complete orthonormal system for $L^2(\mathbb{R})$, made of joint eigenfunctions of the operators (7); in fact,

$$\begin{aligned} d\pi_{\eta,\mu}(L)\tilde{h}_{\eta,n} &= (|\eta|(2n+1)+\mu^2)\tilde{h}_{\eta,n},\\ d\pi_{\eta,\mu}(-iY_j)\tilde{h}_{\eta,n} &= \eta_j \tilde{h}_{\eta,n}. \end{aligned} \tag{8}$$

Moreover the corresponding diagonal matrix coefficients $\varphi_{\eta,\mu,n}$ of $\pi_{\eta,\mu}$ are given by

$$\begin{aligned} \varphi_{\eta,\mu,n}(x,y) &= \langle \pi_{\eta,\mu}(x,y)\tilde{h}_{\eta,n}, \tilde{h}_{\eta,n}\rangle \\ &= e^{i\langle\eta,y\rangle} e^{i\mu x^\eta_\parallel} |\eta|^{1/2} \int_{\mathbb{R}} e^{i|\eta|ux^\eta_2}\, h_n(|\eta|^{1/2}(u+x^\eta_1/2))\, h_n(|\eta|^{1/2}(u-x^\eta_1/2))\,du. \end{aligned}$$

The last integral is essentially the Fourier-Wigner transform of the pair (h_n,h_n), whose Fourier transform has a particularly simple expression (cf. [9, formula (1.90)]); the parity of the Hermite functions then yields

$$\begin{aligned} \varphi_{\eta,\mu,n}(x,y) = e^{i\langle\eta,y\rangle} e^{i\mu x^\eta_\parallel} \frac{(-1)^n}{\pi|\eta|} &\int_{\mathbb{R}^2} e^{iv_2x^\eta_2} e^{iv_1x^\eta_1} \\ &\times \int_{\mathbb{R}} e^{-it(2v_1/|\eta|^{1/2})}\, h_n(t+v_2/|\eta|^{1/2})\, h_n(t-v_2/|\eta|^{1/2})\,dt\,dv, \end{aligned}$$

that is,

$$\varphi_{\eta,\mu,n}(x,y) = \frac{1}{2\pi|\eta|} e^{i\langle\eta,y\rangle} e^{i\mu x^\eta_\parallel} \int_{\mathbb{R}^2} e^{iv_1x^\eta_1} e^{iv_2x^\eta_2} \mathcal{L}^{(0)}_n(|v|^2/|\eta|)\,dv \tag{9}$$

(see [21, Theorem 1.3.4] or [9, Theorem 1.104]).

Note that $\mathcal{K}_{G(L,\mathbf{Y})} \in \mathcal{S}(N_{3,2})$ since $G \in \mathcal{S}(\mathbb{R}^4)$ (see [2, Theorem 5.2] or [12, §4.2]). Moreover

$$\pi_{\eta,\mu}(\mathcal{K}_{G(L,\mathbf{Y})})\tilde{h}_{\eta,n} = G(|\eta|(2n+1)+\mu^2,\eta)\tilde{h}_{\eta,n}$$

by (8) and [18, Proposition 1.1], hence

$$\langle \pi_{\eta,\mu}(x,y)\pi_{\eta,\mu}(\mathcal{K}_{G(L,\mathbf{Y})})\tilde{h}_{\eta,n}, \tilde{h}_{\eta,n}\rangle = m(n,\mu,\eta)\,\varphi_{\eta,\mu,n}(x,y).$$

Therefore, by (6) and (9),

$$\begin{aligned} &\mathcal{K}_{G(L,\mathbf{Y})}(x,y) \\ &\quad= (2\pi)^{-5} \int_{\mathbb{R}^3\setminus\{0\}} \int_{\mathbb{R}} \sum_{n\in\mathbb{N}} m(n,\mu,\eta)\,\varphi_{\eta,\mu,n}(x,y)\,|\eta|\,d\mu\,d\eta \\ &\quad= (2\pi)^{-6} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \sum_{n\in\mathbb{N}} m(n,\xi_3,\eta)\, e^{i\langle\eta,y\rangle} e^{i\langle\xi,(x^\eta_1,x^\eta_2,x^\eta_\parallel)\rangle} \mathcal{L}^{(0)}_n((\xi_1^2+\xi_2^2)/|\eta|)\,d\xi\,d\eta. \end{aligned}$$

The conclusion follows by a change of variable in the inner integral. □

4. Weighted estimates

For convenience, set $\mathcal{L}^{(k)}_n = 0$ for all $n < 0$. The following identities are easily obtained from the properties of Laguerre polynomials (see, e.g., [6, §10.12]).

Lemma 4. *For all $k, n, n' \in \mathbb{N}$ and $t \in \mathbb{R}$,*

$$\mathcal{L}^{(k)}_n(t) = \mathcal{L}^{(k+1)}_{n-1}(t) + \mathcal{L}^{(k+1)}_n(t), \tag{10}$$

$$\frac{d}{dt}\mathcal{L}^{(k)}_n(t) = \mathcal{L}^{(k+1)}_{n-1}(t) - \mathcal{L}^{(k+1)}_n(t), \tag{11}$$

$$\int_0^\infty \mathcal{L}^{(k)}_n(t)\,\mathcal{L}^{(k)}_{n'}(t)\,t^k\,dt = \begin{cases} \frac{(n+k)!}{2^{k-1}n!} & \textit{if } n=n', \\ 0 & \textit{otherwise}. \end{cases} \tag{12}$$

We introduce some operators on functions $f : \mathbb{N} \times \mathbb{R} \times \mathbb{R}^3 \to \mathbb{C}$:

$$\begin{aligned} \tau f(n,\mu,\eta) &= f(n+1,\mu,\eta), \\ \delta f(n,\mu,\eta) &= f(n+1,\mu,\eta) - f(n,\mu,\eta), \\ \partial_\mu f(n,\mu,\eta) &= \frac{\partial}{\partial \mu} f(n,\mu,\eta), \\ \partial_\eta^\alpha f(n,\mu,\eta) &= \left(\frac{\partial}{\partial \eta}\right)^\alpha f(n,\mu,\eta), \end{aligned}$$

for all $\alpha \in \mathbb{N}^3$. For all multiindices $\alpha \in \mathbb{N}^3$, we denote by $|\alpha|$ its length $\alpha_1 + \alpha_2 + \alpha_3$. We set moreover $\langle t \rangle = 2|t| + 1$ for all $t \in \mathbb{R}$.

Note that, for all compactly supported $f : \mathbb{N} \times \mathbb{R} \times \mathbb{R}^3 \to \mathbb{C}$, $\tau^l f$ is null for all sufficiently large $l \in \mathbb{N}$; hence the operator $1 + \tau$, when restricted to the set of compactly supported functions, is invertible, with inverse given by

$$(1+\tau)^{-1} f = \sum_{l \in \mathbb{N}} (-1)^l \tau^l f,$$

and therefore the operator $(1+\tau)^q$ is well-defined for all $q \in \mathbb{Z}$.

Proposition 5. *Let $G : \mathbb{R}^4 \to \mathbb{C}$ be smooth and compactly supported in $\mathbb{R} \times (\mathbb{R}^3 \setminus \{0\})$, and let $m(n,\mu,\eta)$ be defined by* (5)*. For all $\alpha \in \mathbb{N}^3$,*

$$\begin{aligned} &\int_{N_{3,2}} |y^\alpha \, \mathcal{K}_{G(L,\mathbf{Y})}(x,y)|^2 \, dx \, dy \\ &\qquad \leq C_\alpha \sum_{\iota \in I_\alpha} \sum_{n \in \mathbb{N}} \int_{\mathbb{R}^3} \int_{\mathbb{R}} |\partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota| - k_\iota} m(n,\mu,\eta)|^2 \\ &\qquad\qquad \times \mu^{2b_\iota} \, |\eta|^{2|\gamma^\iota| - 2|\alpha| - 2k_\iota + |\beta^\iota| + 1} \langle n \rangle^{|\beta^\iota|} \, d\mu \, d\eta, \end{aligned} \tag{13}$$

where I_α is a finite set and, for all $\iota \in I_\alpha$,

- $\gamma^\iota \in \mathbb{N}^3$, $l_\iota, k_\iota \in \mathbb{N}$, $\gamma^\iota \leq \alpha$, $\min\{1, |\alpha|\} \leq |\gamma^\iota| + l_\iota + k_\iota \leq |\alpha|$,
- $b_\iota \in \mathbb{N}$, $\beta^\iota \in \mathbb{N}^3$, $b_\iota + |\beta^\iota| = l_\iota + 2k_\iota$, $|\gamma^\iota| + l_\iota + b_\iota \leq |\alpha|$.

Proof. Proposition 3 and integration by parts allow us to write

$$\begin{aligned} &y^\alpha \, \mathcal{K}_{G(L,\mathbf{Y})}(x,y) \\ &\quad = \frac{i^{|\alpha|}}{(2\pi)^6} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \left[\left(\frac{\partial}{\partial \eta}\right)^\alpha \sum_{n \in \mathbb{N}} m(n, \xi_\parallel^\eta, \eta) \, \mathcal{L}_n^{(0)}(|\xi_\perp^\eta|^2/|\eta|) \right] e^{i\langle \xi, x \rangle} \, e^{i \langle \eta, y \rangle} \, d\xi \, d\eta. \end{aligned} \tag{14}$$

From the definition of $\xi_\parallel^\eta$ and $\xi_\perp^\eta$, the following identities are not difficult to obtain:

$$\begin{gathered} \frac{\partial}{\partial \eta_j} \xi_\parallel^\eta = (\xi_\perp^\eta)_j \frac{1}{|\eta|}, \qquad \frac{\partial}{\partial \eta_j} (\xi_\perp^\eta)_k = -\xi_\parallel^\eta \frac{\partial}{\partial \eta_j} \frac{\eta_k}{|\eta|} - (\xi_\perp^\eta)_j \frac{\eta_k}{|\eta|^2}, \\ \frac{\partial}{\partial \eta_j} \frac{|\xi_\perp^\eta|^2}{|\eta|} = -\xi_\parallel^\eta (\xi_\perp^\eta)_j \frac{2}{|\eta|^2} - |\xi_\perp^\eta|^2 \frac{\eta_j}{|\eta|^3}. \end{gathered} \tag{15}$$

The multiindex notation will also be used as follows:

$$(\xi_\perp^\eta)^\beta = (\xi_\perp^\eta)_1^{\beta_1} (\xi_\perp^\eta)_2^{\beta_2} (\xi_\perp^\eta)_3^{\beta_3}$$

for all $\xi, \eta \in \mathbb{R}$, with $\eta \neq 0$, and all $\beta \in \mathbb{N}^3$; consequently

$$|\xi_\perp^\eta|^2 = (\xi_\perp^\eta)^{(2,0,0)} + (\xi_\perp^\eta)^{(0,2,0)} + (\xi_\perp^\eta)^{(0,0,2)}.$$

Via these identities, one can prove inductively that, for all $\alpha \in \mathbb{N}^3$,

$$\begin{aligned}(16)\qquad \left(\frac{\partial}{\partial \eta}\right)^{\alpha} &\sum_{n \in \mathbb{N}} m(n, \xi^\eta_\parallel, \eta)\, \mathcal{L}_n^{(0)}(|\xi^\eta_\perp|^2/|\eta|) \\ &= \sum_{\iota \in I_\alpha} \sum_{n \in \mathbb{N}} \partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} m(n, \xi^\eta_\parallel, \eta)\, (\xi^\eta_\parallel)^{b_\iota}\, (\xi^\eta_\perp)^{\beta^\iota}\, \Theta_\iota(\eta)\, \mathcal{L}_n^{(k_\iota)}(|\xi^\eta_\perp|^2/|\eta|),\end{aligned}$$

where I_α, γ^ι, l_ι, k_ι, b_ι, β^ι are as in the statement above, while $\Theta_\iota : \mathbb{R}^3 \setminus \{0\} \to \mathbb{R}$ is smooth and homogeneous of degree $|\gamma^\iota| - |\alpha| - k_\iota$. For the inductive step, one employs Leibniz' rule, and when a derivative hits a Laguerre function, the identity (11) together with summation by parts is used.

Note that, for all compactly supported $f : \mathbb{N} \times \mathbb{R} \times \mathbb{R}^3 \to \mathbb{C}$,

$$\sum_{n \in \mathbb{N}} f(n, \mu, \eta)\, \mathcal{L}_n^{(k)}(t) = \sum_{n \in \mathbb{N}} (1 + \tau) f(n, \mu, \eta)\, \mathcal{L}_n^{(k+1)}(t),$$

by (10). Since $1 + \tau$ is invertible, simple manipulations and iteration yield the more general identity

$$\sum_{n \in \mathbb{N}} f(n, \mu, \eta)\, \mathcal{L}_n^{(k)}(t) = \sum_{n \in \mathbb{N}} (1 + \tau)^{k'-k} f(n, \mu, \eta)\, \mathcal{L}_n^{(k')}(t),$$

for all $k, k' \in \mathbb{N}$. This formula allows us to adjust in (16) the type of the Laguerre functions to the exponent of $\xi_\perp$, and to obtain that

$$\begin{aligned}&\left(\frac{\partial}{\partial \eta}\right)^{\alpha} \sum_{n \in \mathbb{N}} m(n, \xi^\eta_\parallel, \eta)\, \mathcal{L}_n^{(0)}(|\xi^\eta_\perp|^2/|\eta|) \\ &= \sum_{\iota \in I_\alpha} \sum_{n \in \mathbb{N}} \partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota| - k_\iota} m(n, \xi^\eta_\parallel, \eta)\, (\xi^\eta_\parallel)^{b_\iota}\, (\xi^\eta_\perp)^{\beta^\iota}\, \Theta_\iota(\eta)\, \mathcal{L}_n^{(|\beta^\iota|)}(|\xi^\eta_\perp|^2/|\eta|),\end{aligned}$$

By plugging this identity into (14) and exploiting Plancherel's formula for the Fourier transform, the finiteness of I_α and the triangular inequality, we get that

$$\begin{aligned}&\int_{N_{3,2}} |y^\alpha\, \mathcal{K}_{G(L,\mathbf{Y})}(x, y)|^2 \,dx\,dy \\ &\quad \leq C_\alpha \sum_{\iota \in I_\alpha} \int_{\mathbb{R}^3} \int_{\mathbb{R}} \int_{\mathbb{R}^2} \left| \sum_{n \in \mathbb{N}} \partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota| - k_\iota} m(n, \mu, \eta)\, \mathcal{L}_n^{(|\beta^\iota|)}(|\zeta|^2/|\eta|) \right|^2 \\ &\qquad\qquad\qquad \times \mu^{2b_\iota}\, |\zeta|^{2|\beta^\iota|}\, |\eta|^{2|\gamma^\iota| - 2|\alpha| - 2k_\iota} \,d\zeta\,d\mu\,d\eta\end{aligned}$$

A passage to polar coordinates in the ζ-integral and a rescaling then give that

$$\begin{aligned}&\int_{N_{3,2}} |y^\alpha\, \mathcal{K}_{G(L,\mathbf{Y})}(x, y)|^2 \,dx\,dy \\ &\quad \leq C_\alpha \sum_{\iota \in I_\alpha} \int_{\mathbb{R}^3} \int_{\mathbb{R}} \int_0^\infty \left| \sum_{n \in \mathbb{N}} \partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota| - k_\iota} m(n, \mu, \eta)\, \mathcal{L}_n^{(|\beta^\iota|)}(s) \right|^2 s^{|\beta^\iota|} \,ds \\ &\qquad\qquad\qquad \times \mu^{2b_\iota}\, |\eta|^{2|\gamma^\iota| - 2|\alpha| - 2k_\iota + |\beta^\iota| + 1} \,d\mu\,d\eta,\end{aligned}$$

and the conclusion follows by applying the orthogonality relations (12) for the Laguerre functions to the inner integral. □

Note that $\tau f(\cdot, \mu, \eta)$, $\delta f(\cdot, \mu, \eta)$ depend only on $f(\cdot, \mu, \eta)$; in other words, τ and δ can be considered as operators on functions $\mathbb{N} \to \mathbb{C}$. The next lemma will be useful in converting finite differences into continuous derivatives.

Lemma 6. *Let* $f : \mathbb{N} \to \mathbb{C}$ *have a smooth extension* $\tilde{f} : [0, \infty[\to \mathbb{C}$*, and let* $k \in \mathbb{N}$*. Then*

$$\delta^k f(n) = \int_{J_k} \tilde{f}^{(k)}(n+s)\, d\nu_k(s)$$

for all $n \in \mathbb{N}$*, where* $J_k = [0,k]$ *and* ν_k *is a Borel probability measure on* J_k*. In particular*

$$|\delta^k f(n)|^2 \leq \int_{J_k} |\tilde{f}^{(k)}(n+s)|^2\, d\nu_k(s)$$

for all $n \in \mathbb{N}$*.*

Proof. Iterated application of the fundamental theorem of integral calculus gives

$$\delta^k f(n) = \int_{[0,1]^k} \tilde{f}^{(k)}(n + s_1 + \dots + s_k)\, ds.$$

The conclusion follows by taking as ν_k the push-forward of the uniform distribution on $[0,1]^k$ via the map $(s_1, \dots, s_k) \mapsto s_1 + \dots + s_k$, and by Hölder's inequality. ☐

We give now a simplified version of the right-hand side of (13), in the case where we restrict to the functional calculus for the sublaplacian L alone. In order to avoid divergent series, however, it is convenient at first to truncate the multiplier along the spectrum of $\mathbf{Y}$.

Lemma 7. *Let* $\chi \in C_c^\infty(\mathbb{R})$ *be supported in* $[1/2, 2]$*,* $K \subseteq]0, \infty[$ *be compact and* $M \in]0, \infty[$*. If* $F : \mathbb{R} \to \mathbb{C}$ *is smooth and supported in* K*, and* $F_M : \mathbb{R} \times \mathbb{R}^3 \to \mathbb{C}$ *is given by*

$$F_M(\lambda, \eta) = F(\lambda)\, \chi(|\eta|/M),$$

then, for all $r \in [0, \infty[$*,*

$$\int_{N_{3,2}} ||y|^r\, \mathcal{K}_{F_M(L,\mathbf{Y})}(x,y)|^2\, dx\, dy \leq C_{K,\chi,r}\, M^{3-2r} \|F\|^2_{W_2^r}.$$

Proof. We may restrict to the case $r \in \mathbb{N}$, the other cases being recovered a posteriori by interpolation. Hence we need to prove that

$$\int_{N_{3,2}} |y^\alpha\, \mathcal{K}_{F_M(L,\mathbf{Y})}(x,y)|^2\, dx\, dy \leq C_{K,\chi,\alpha}\, M^{3-2|\alpha|} \|F\|^2_{W_2^{|\alpha|}} \tag{17}$$

for all $\alpha \in \mathbb{N}^3$. On the other hand, if

$$m(n, \mu, \eta) = F(|\eta| \langle n \rangle + \mu^2)\, \chi(|\eta|/M),$$

then the left-hand side of (17) can be majorized by (13), and we are reduced to proving

$$\begin{aligned} \sum_{n \in \mathbb{N}} \int_{\mathbb{R}^3} \int_{\mathbb{R}} |\partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota| - k_\iota} m(n,\mu,\eta)|^2\, \mu^{2b_\iota}\, |\eta|^{2|\gamma^\iota| - 2|\alpha| - 2k_\iota + |\beta^\iota| + 1} \\ \times \langle n \rangle^{|\beta^\iota|}\, d\mu\, d\eta \leq C_{K,\chi,\alpha}\, M^{3-2|\alpha|} \|F\|^2_{W_2^{|\alpha|}} \end{aligned} \tag{18}$$

for all $\iota \in I_\alpha$.

Consider first the case $|\beta^\iota| \geq k_\iota$. A smooth extension $\tilde{m} : \mathbb{R} \times \mathbb{R} \times \mathbb{R}^3 \to \mathbb{C}$ of m is defined by

$$\tilde{m}(t, \mu, \eta) = F(|\eta|(2t+1) + \mu^2)\, \chi(|\eta|/M).$$

Then, by Lemma 6,

$$\begin{aligned} &\partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota| - k_\iota} m(n,\mu,\eta) \\ &\qquad = \sum_{j=0}^{|\beta^\iota| - k_\iota} \binom{|\beta^\iota| - k_\iota}{j} \int_{J_\iota} \partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \partial_t^{k_\iota} \tilde{m}(n+j+s, \mu, \eta)\, d\nu_\iota(s), \end{aligned}$$

where $J_\iota = [0, k_\iota]$ and ν_ι is a suitable probability measure on J_ι; consequently (18) will be proved if we show that

$$\begin{aligned}(19)\quad \sum_{n\in\mathbb{N}} \int_{\mathbb{R}^3}\int_{\mathbb{R}} |\partial_\eta^{\gamma^\iota}\partial_\mu^{l_\iota}\partial_t^{k_\iota}\tilde{m}(n+s,\mu,\eta)|^2\, \mu^{2b_\iota}\, |\eta|^{2|\gamma^\iota|-2|\alpha|-2k_\iota+|\beta^\iota|+1} \\ \times \langle n\rangle^{|\beta^\iota|}\, d\mu\, d\eta \le C_{K,\chi,\alpha}\, M^{3-2|\alpha|} \|F\|^2_{W_2^{|\alpha|}}\end{aligned}$$

for all $s \in [0, |\beta^\iota|]$. On the other hand, it is easily proved inductively that

$$\begin{aligned}&\partial_\eta^{\gamma^\iota}\partial_\mu^{l_\iota}\partial_t^{k_\iota}\tilde{m}(t,\mu,\eta) \\ &= \sum_{r=\lceil l_\iota/2\rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} \sum_{q=0}^{|\gamma^\iota|-v} \Psi_{i,v,q}(\eta)\, \langle t\rangle^v \mu^{2r-l_\iota} M^{-q} F^{(k_\iota+v+r)}(|\eta|\langle t\rangle + \mu^2)\, \chi^{(q)}(|\eta|/M)\end{aligned}$$

for all $t \geq 0$, where $\Psi_{i,q,v} : \mathbb{R}^3 \setminus \{0\} \to \mathbb{R}$ is smooth and homogeneous of degree $k_\iota + v + q - |\gamma^\iota|$; hence

$$\begin{aligned}(20)\quad |\partial_\eta^{\gamma^\iota}\partial_\mu^{l_\iota}\partial_t^{k_\iota}\tilde{m}(t,\mu,\eta)|^2 \le C_{\chi,\alpha} \sum_{r=\lceil l_\iota/2\rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} M^{2k_\iota+2v-2|\gamma^\iota|} \langle t\rangle^{2v} \mu^{4r-2l_\iota} \\ \times |F^{(k_\iota+v+r)}(|\eta|\langle t\rangle + \mu^2)|^2\, \tilde{\chi}(|\eta|/M),\end{aligned}$$

where $\tilde{\chi}$ is the characteristic function of $[1/2, 2]$, and we are using the fact that $|\eta| \sim M$ in the region where $\tilde{\chi}(|\eta|/M) \neq 0$. Consequently the left-hand side of (19) is majorized by

$$\begin{aligned}&C_{\chi,\alpha} \sum_{r=\lceil l_\iota/2\rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} M^{2v-2|\alpha|+|\beta^\iota|+1} \sum_{n\in\mathbb{N}} \langle n\rangle^{|\beta^\iota|} \langle n+s\rangle^{2v} \\ &\qquad \times \int_{\mathbb{R}^3}\int_{\mathbb{R}} |F^{(k_\iota+v+r)}(|\eta|\langle n+s\rangle + \mu^2)|^2\, \mu^{2b_\iota+4r-2l_\iota}\, \tilde{\chi}(|\eta|/M)\, d\mu\, d\eta \\ &\le C_{\chi,\alpha} \sum_{r=\lceil l_\iota/2\rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} M^{2v-2|\alpha|+|\beta^\iota|+3} \sum_{n\in\mathbb{N}} \langle n+s\rangle^{|\beta^\iota|+2v} \\ &\qquad \times \int_0^\infty \int_0^\infty |F^{(k_\iota+v+r)}(\rho\langle n+s\rangle + \mu^2)|^2\, \mu^{2b_\iota+4r-2l_\iota}\, \tilde{\chi}(\rho/M)\, d\mu\, d\rho \\ &\le C_{\chi,\alpha} \sum_{r=\lceil l_\iota/2\rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} M^{2v-2|\alpha|+|\beta^\iota|+3} \int_0^\infty \int_0^\infty |F^{(k_\iota+v+r)}(\rho + \mu^2)|^2 \\ &\qquad \times \mu^{2b_\iota+4r-2l_\iota} \sum_{n\in\mathbb{N}} \langle n+s\rangle^{|\beta^\iota|+2v-1} \tilde{\chi}(\rho/(\langle n+s\rangle M))\, d\mu\, d\rho,\end{aligned}$$

by passing to polar coordinates and rescaling. The last sum in n is easily controlled by $(\rho/M)^{|\beta^\iota|+2v}$, hence the left-hand side of (19) is majorized by

$$\begin{aligned}&C_{\chi,\alpha}\, M^{3-2|\alpha|} \sum_{r=\lceil l_\iota/2\rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} \int_0^\infty \int_0^\infty |F^{(k_\iota+v+r)}(\rho+\mu^2)|^2 \mu^{2b_\iota+4r-2l_\iota}\, \rho^{|\beta^\iota|+2v}\, d\mu\, d\rho \\ &\qquad \le C_{K,\chi,\alpha}\, M^{3-2|\alpha|} \sum_{r=\lceil l_\iota/2\rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} \sup_{u\in[0,\max K]} \int_0^\infty |F^{(k_\iota+v+r)}(\rho+u)|^2\, d\rho,\end{aligned}$$

because $2b_\iota + 4r - 2l_\iota \geq 0$ and $|\beta^\iota| + 2v \geq 0$ if r and v are in the range of summation, and $\operatorname{supp} F \subseteq K$. Since moreover $k_\iota + v + r \leq k_\iota + \gamma^\iota + l_\iota \leq |\alpha|$, the last integral is dominated by $\|F\|_{W_2^{|\alpha|}}$ uniformly in r, v, u, and (19) follows.

Consider now the case $|\beta^\iota| < k_\iota$. Via the identity

$$(1+\tau)^{-1} = (1-\tau)(1-\tau^2)^{-1} = -\delta(1-\tau^2)^{-1} = -\delta \sum_{j=0}^\infty \tau^{2j},$$

together with Lemma 6, we obtain that

$$\begin{aligned} (21) \quad & \partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota|-k_\iota} m(n,\mu,\eta) \\ & = (-1)^{k_\iota - |\beta^\iota|} \sum_{j=0}^\infty \tbinom{j+k_\iota-|\beta^\iota|-1}{k_\iota-|\beta^\iota|-1} \int_{J_\iota} \partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \partial_t^{2k_\iota-|\beta^\iota|} \tilde{m}(n+2j+s,\mu,\eta)\, d\nu_\iota(s), \end{aligned}$$

where $J_\iota = [0, 2k_\iota - |\beta^\iota|]$ and ν_ι is a suitable probability measure on J_ι. Note that, because of the assumptions on the supports of F and χ, the sum on j in the right-hand side of (21) is a finite sum, that is, the j-th summand is nonzero only if $\langle n+2j \rangle \leq 2M^{-1} \max K$; consequently, by applying the Cauchy-Schwarz inequality to the sum in j, and by (20),

$$\begin{aligned} & |\partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \delta^{k_\iota} (1+\tau)^{|\beta^\iota|-k_\iota} m(n,\mu,\eta)|^2 \\ & \quad \leq C_{K,\alpha}\, M^{1+2|\beta^\iota|-2k_\iota} \sum_{j=0}^\infty \int_{J_\iota} |\partial_\eta^{\gamma^\iota} \partial_\mu^{l_\iota} \partial_t^{2k_\iota-|\beta^\iota|} \tilde{m}(n+2j+s,\mu,\eta)|^2 \, d\nu_\iota(s) \\ & \quad \leq C_{K,\chi,\alpha} \sum_{r=\lceil l_\iota/2 \rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} M^{1+2k_\iota+2v-2|\gamma^\iota|} \sum_{j=0}^\infty \int_{J_\iota} \langle n+2j+s \rangle^{2v} \mu^{4r-2l_\iota} \\ & \qquad\qquad \times |F^{(2k_\iota-|\beta^\iota|+v+r)}(|\eta|\langle n+2j+s\rangle + \mu^2)|^2\, \tilde{\chi}(|\eta|/M)\, d\nu_\iota(s). \end{aligned}$$

Remember that $|\eta| \sim M$ in the region where $\tilde{\chi}(|\eta|/M) \neq 0$. Hence the left-hand side of (18) is majorized by

$$\begin{aligned} & C_{K,\chi,\alpha} \sum_{r=\lceil l_\iota/2 \rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} \int_{J_\iota} \sum_{n\in\mathbb{N}} \sum_{j\in\mathbb{N}} \langle n+2j+s\rangle^{2v} \langle n \rangle^{|\beta^\iota|} \int_{\mathbb{R}^3} \int_{\mathbb{R}} M^{2+2v-2|\alpha|+|\beta^\iota|} \\ & \quad \times \mu^{2b_\iota+4r-2l_\iota}\, |F^{(2k_\iota-|\beta^\iota|+v+r)}(|\eta|\langle n+2j+s\rangle+\mu^2)|^2\, \tilde{\chi}(|\eta|/M)\, d\mu\, d\eta\, d\nu_\iota(s) \\ & \leq C_{K,\chi,\alpha} \sum_{r=\lceil l_\iota/2 \rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} \int_{J_\iota} \sum_{n\in\mathbb{N}} \sum_{j\in\mathbb{N}} \langle n+2j+s\rangle^{2v+|\beta^\iota|} \int_0^\infty \int_0^\infty M^{4+2v-2|\alpha|+|\beta^\iota|} \\ & \quad \times \mu^{2b_\iota+4r-2l_\iota}\, |F^{(2k_\iota-|\beta^\iota|+v+r)}(\rho\langle n+2j+s\rangle+\mu^2)|^2\, \tilde{\chi}(\rho/M)\, d\mu\, d\rho\, d\nu_\iota(s) \\ & \leq C_{K,\chi,\alpha} \sum_{r=\lceil l_\iota/2 \rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} M^{4+2v-2|\alpha|+|\beta^\iota|} \int_0^\infty \int_0^\infty |F^{(2k_\iota-|\beta^\iota|+v+r)}(\rho+\mu^2)|^2 \\ & \quad \times \mu^{2b_\iota+4r-2l_\iota} \int_{J_\iota} \sum_{(n,j)\in\mathbb{N}^2} \langle n+2j+s\rangle^{2v+|\beta^\iota|-1} \tilde{\chi}(\rho/(\langle n+2j+s\rangle M))\, d\nu_\iota(s)\, d\mu\, d\rho, \end{aligned}$$

by passing to polar coordinates and rescaling. The sum in (n,j) is dominated by $(\rho/M)^{2v+|\beta^\iota|+1}$, uniformly in $s \in J_\iota$, and moreover $\operatorname{supp} F \subseteq K$. Therefore the left-hand side of (18) is majorized by

$$C_{K,\chi,\alpha}\, M^{3-2|\alpha|} \sum_{r=\lceil l_\iota/2 \rceil}^{l_\iota} \sum_{v=0}^{|\gamma^\iota|} \sup_{u\in[0,\max K]} \int_0^\infty |F^{(2k_\iota-|\beta^\iota|+v+r)}(\rho+u)|^2\, d\rho.$$

On the other hand, $b_\iota + |\beta^\iota| = l_\iota + 2k_\iota$, hence $2k_\iota - |\beta^\iota| + v + r \leq 2k_\iota - |\beta^\iota| + |\gamma^\iota| + l_\iota = b_\iota + |\gamma^\iota| \leq |\alpha|$ if r and v are in the range of summation, therefore the last integral is dominated by $\|F\|_{W_2^{|\alpha|}}$ uniformly in r, v, u, and (18) follows. □

Proposition 8. *Let* $F : \mathbb{R} \to \mathbb{C}$ *be smooth and such that* $\operatorname{supp} F \subseteq K$ *for some compact set* $K \subseteq]0,\infty[$*. For all* $r \in [0, 3/2[$*,*

$$\int_{N_{3,2}} \left|(1+|y|)^r \, \mathcal{K}_{F(L)}(x,y)\right|^2 \, dx \, dy \leq C_{K,r} \|F\|^2_{W_2^r}.$$

Proof. Take $\chi \in C_c^\infty(]0,\infty[)$ such that $\operatorname{supp} \chi \subseteq [1/2, 2]$ and $\sum_{k\in\mathbb{Z}} \chi(2^{-k}t) = 1$ for all $t \in]0,\infty[$. Note that, if (λ,η) belongs to the joint spectrum of $L, \mathbf{Y}$, then $|\eta| \leq \lambda$. Therefore, if $k_K \in \mathbb{Z}$ is sufficiently large so that $2^{k_K - 1} > \max K$, and if F_M is defined for all $M \in]0,\infty[$ as in Lemma 7, then

$$F(L) = \sum_{k\in\mathbb{Z},\, k\leq k_K} F_{2^k}(L, \mathbf{Y})$$

(with convergence in the strong sense). Hence an estimate for $\mathcal{K}_{F(L)}$ can be obtained, via Minkowski's inequality, by summing the corresponding estimates for $\mathcal{K}_{F_{2^k}}(L, \mathbf{Y})$ given by Lemma 7. If $r < 3/2$, then the series $\sum_{k\leq k_K} (2^k)^{3/2-r}$ converges, thus

$$\int_{N_{3,2}} \left||y|^r \, \mathcal{K}_{F(L)}(x,y)\right|^2 \, dx \, dy \leq C_{K,r} \|F\|^2_{W_2^r}.$$

The conclusion follows by combining the last inequality with the corresponding one for $r = 0$. □

Recall that $|\cdot|_\delta$ denotes a δ_t-homogeneous norm on $N_{3,2}$, thus $|(x,y)|_\delta \sim |x| + |y|^{1/2}$. Interpolation then allows us to improve the standard weighed estimate for a homogeneous sublaplacian on a stratified group.

Proposition 9. *Let* $F : \mathbb{R} \to \mathbb{C}$ *be smooth and such that* $\operatorname{supp} F \subseteq K$ *for some compact set* $K \subseteq]0,\infty[$*. For all* $r \in [0, 3/2[$*,* $\alpha \geq 0$ *and* $\beta > \alpha + r$*,*

$$\int_{N_{3,2}} \left|(1+|(x,y)|_\delta)^\alpha \, (1+|y|)^r \, \mathcal{K}_{F(L)}(x,y)\right|^2 \, dx \, dy \leq C_{K,\alpha,\beta,r} \|F\|^2_{W_2^\beta}. \tag{22}$$

Proof. Note that $1 + |y| \leq C(1 + |(x,y)|_\delta)^2$. Hence, in the case $\alpha \geq 0$, $\beta > \alpha + 2r$, the inequality (22) follows by the standard estimate [16, Lemma 1.2]. On the other hand, if $\alpha = 0$ and $\beta \geq r$, then (22) is given by Proposition 8. The full range of α and β is then obtained by interpolation (cf. the proof of [16, Lemma 1.2]). □

We can finally prove the fundamental L^1-estimate, and consequently Theorem 1.

Proof of Proposition 2. Take $r \in]9/2 - s, 3/2[$. Then $s - r > 3/2 + 3 - 2r$, hence we can find $\alpha_1 > 3/2$ and $\alpha_2 > 3 - 2r$ such that $s - r > \alpha_1 + \alpha_2$. Therefore, by Proposition 9 and Hölder's inequality,

$$\| \mathcal{K}_{F(L)} \|_1^2 \leq C_{k,s} \|F\|^2_{W_2^s} \int_{N_{3,2}} (1+|(x,y)|_\delta)^{-2\alpha_1 - 2\alpha_2} \, (1+|y|)^{-2r} \, dx \, dy.$$

The integral on the right-hand side is finite, because $2\alpha_1 > 3$, $\alpha_2 + 2r > 3$, and

$$(1+|(x,y)|_\delta)^{-2\alpha_1-2\alpha_2} \, (1+|y|)^{-2r} \leq C_s (1+|x|)^{-2\alpha_1} \, (1+|y|)^{-\alpha_2-2r},$$

and we are done. □

References

[1] F. Astengo, M. Cowling, B. Di Blasio, and M. Sundari, *Hardy's uncertainty principle on certain Lie groups*, J. London Math. Soc. (2) **62** (2000), no. 2, 461–472.

[2] F. Astengo, B. Di Blasio, and F. Ricci, *Gelfand pairs on the Heisenberg group and Schwartz functions*, J. Funct. Anal. **256** (2009), no. 5, 1565–1587.

[3] M. Christ, *L^p bounds for spectral multipliers on nilpotent groups*, Trans. Amer. Math. Soc. **328** (1991), no. 1, 73–81.

[4] L. De Michele and G. Mauceri, *L^p multipliers on the Heisenberg group*, Michigan Math. J. **26** (1979), no. 3, 361–371.

[5] ———, *H^p multipliers on stratified groups*, Ann. Mat. Pura Appl. (4) **148** (1987), 353–366.

[6] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher transcendental functions. Vol. II*, Robert E. Krieger Publishing Co. Inc., Melbourne, Fla., 1981.

[7] V. Fischer and F. Ricci, *Gelfand transforms of* SO(3)*-invariant Schwartz functions on the free group $N_{3,2}$*, Ann. Inst. Fourier (Grenoble) **59** (2009), no. 6, 2143–2168.

[8] G. B. Folland and E. M. Stein, *Hardy spaces on homogeneous groups*, Mathematical Notes, vol. 28, Princeton University Press, Princeton, N.J., 1982.

[9] G. B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies, vol. 122, Princeton University Press, Princeton, NJ, 1989.

[10] W. Hebisch, *Multiplier theorem on generalized Heisenberg groups*, Colloq. Math. **65** (1993), no. 2, 231–239.

[11] W. Hebisch and J. Zienkiewicz, *Multiplier theorem on generalized Heisenberg groups. II*, Colloq. Math. **69** (1995), no. 1, 29–36.

[12] A. Martini, *Algebras of differential operators on Lie groups and spectral multipliers*, Tesi di perfezionamento (PhD thesis), Scuola Normale Superiore, Pisa, 2010, `arXiv:1007.1119`.

[13] A. Martini, *Spectral theory for commutative algebras of differential operators on Lie groups*, J. Funct. Anal. **260** (2011), no. 9, 2767–2814.

[14] ———, *Analysis of joint spectral multipliers on Lie groups of polynomial growth*, Ann. Inst. Fourier (Grenoble) **62** (2012), no. 4, 1215–1263.

[15] A. Martini and D. Müller, *A sharp multiplier theorem for Grushin operators in arbitrary dimensions*, (2012), `arXiv:1210.3564`.

[16] G. Mauceri and S. Meda, *Vector-valued multipliers on stratified groups*, Rev. Mat. Iberoamericana **6** (1990), no. 3-4, 141–154.

[17] D. Müller and E. M. Stein, *On spectral multipliers for Heisenberg and related groups*, J. Math. Pures Appl. (9) **73** (1994), no. 4, 413–440.

[18] D. Müller, *A restriction theorem for the Heisenberg group*, Ann. of Math. (2) **131** (1990), no. 3, 567–587.

[19] D. Müller, F. Ricci, and E. M. Stein, *Marcinkiewicz multipliers and multi-parameter structure on Heisenberg (-type) groups. II*, Math. Z. **221** (1996), no. 2, 267–291.

[20] L. P. Rothschild and E. M. Stein, *Hypoelliptic differential operators and nilpotent groups*, Acta Math. **137** (1976), no. 3-4, 247–320.

[21] S. Thangavelu, *Lectures on Hermite and Laguerre expansions*, Mathematical Notes, vol. 42, Princeton University Press, Princeton, NJ, 1993.

ALESSIO MARTINI, MATHEMATISCHES SEMINAR, CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL, LUDEWIG-MEYN-STR. 4, D-24118 KIEL, GERMANY
E-mail address: `martini@math.uni-kiel.de`

DETLEF MÜLLER, MATHEMATISCHES SEMINAR, CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL, LUDEWIG-MEYN-STR. 4, D-24118 KIEL, GERMANY
E-mail address: `mueller@math.uni-kiel.de`